

Oborisy Cinc. p. 114 That of 1 De Bod C. pedrerdanis Kold - ochel. 18453 This species so well diligned by the truste Celyx and the about entre scales bear be for new Days & South Day 14 20 x 3183) At Tyris & . Noi & i Kuntit ly Kitch (388) MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS

O. elegans Boids & Dalaum Sing, N. on.
II.3. 129

for the alpian rights of the Teams

(Dalaum N. 708)


is cravily deliganthe except bythe

papilloss prettiles rose rolat flowers a

the scales been a Cittle surre income


0 1 2 3 4 5 6 7 8 9 10

Copyright reserved



Casoute Abyssiaira Mich. Abyss. II.78 C'mentelle Dearin - H. Mus Vani. Teens the well distinguild by the short thick love of Halyx A the very log to namon lobes of the Corolla, the snall, off biful scales a the veg legg, capilles style, The more flowing the thick calyx as the laft of sales approach it te C. par flow, while the sleader offle en surpens that of Epithy. After way anoth cott sant voulle lober was collect a son Egencery should sent sent





Cascuta planiflora Tenore Sol.

Carle capillació glomentis globosis mellifloris arrite orgestin Horibus sessitibus planague of ides planague of ides training tumidis intumentation tomaine touristics consollar applies s. superatibus; lariais tubem aequitibus accitionsalis s. accitis, planger aproceptes miner tames centibus . Derecti, dence hørjetet. E. pelkatiber, square adpressis su incomis, orati, truveti s. bifits. style skriguente agentrege oppførie kongrøre gleon oranien definites ogløbopen, rapsula river, rissa, corolla manera odge skoolate, semile (aspergtifvermords-) Mostle This Ditt wasnur parts of lumpe Cashing Mostle Africa, the Cananies, and the Clark Leave atthus seite Asi t the Municipalland - Sher externely vanible speries has been a juggle tack believet, esto part attack. to Cerculier, and on our ask some that entirely Tale for with my ettenflet answer the difficulty. I have been obliged to unite a series of former the extreme of which are so willy lifered from our authority a careful study of the even varying for chemitan could have where me not to keep the Texante litt the 190 sperime, before are from through, Aprin e Asi I fine it suposissille to draw the liverity of the subsperies while sex 6 m & spening might be delect apportly to different that they with be used as types of well eshablishe species on the other Mand, official Sifferell to Keep Their species or complex of forms as I would prefer to call it, separt for Nouve allier spiries p some form approach

copyright reserved cm

MISSOURI BOTANICAL GARDEN

to Ceasent brevislore solvich might be comished a short styles form of our plant, others to C. faleatin other again and diffiell to separate fritt alfine form of C. Grillyon, 10 That an species find et, als jances alevort catine orrele of the rubgeau, Synony for The speriet on I view it there are none at Leaver nam itself belong property speaking ong to one of the form, but a leg list frynongen vill be for with the differe July comp. MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS



Casonta Valacetina Obviss. C. Palestir. Bors, Siegn. T. 9. 86. Welf. Chun. III. 118. C. alba var. Briss Riagn., Heldrich School. C. Cretica Tournet. con. 45. C. mirantha Tive o fide sper auct. I - Jussone Hor Siral Syn. II. 887. (hold) non Choisy Mem (1841)
Coopellarie Recibel. M. Crit. V. p. G. (1824) -sira desimpt. This pretty little species the glowner of Nach have de 2 ling i hat the flows /2-1 l. log. tt send 0.5 l. - Samten, is weathy aller to the planiflow to while - e It nam of Calle the authorite subsequely referme it. Bet I believe it to be seffy vel detypiste et their its such with ander fothe speries of the golding the Ciaits of while are is diffull to I cover Townfort den z Hillegent it at the object spening the Compact herby. Then of Sieben rown for Tourful, all Confe the Budi or Cretz, It is public ful åre tt tilledtemmen, sin dy hill good on my by plate That as the Tale Shubby Labretter of those vigory to I have seen thereto for Blacking (Borgo) H. Libanon, Blanche, Garllande atter Heldrich Morra Bong 2. 11 Uncl Creta Siban Raulin and other Green islands Raulin, Seferire Sibant ad for Sirily, Times MISSOURI BOTANICAL copyright reserved


GARDEN

x cutt on frem ten flow of else stat His Whinghow by the flow, beginning to be mostly to party lage, loby bout unte goldenby rante n back top of rosollier lærnie omrellet. Tunid, rule, sund utte large orbille or bruty ovel, somethe Housel, The short be longer the the note langer orang. Lieber Lieberg and Millian Committee of the sound of the state of the colones the state of the s A STANDARD OF THE STANDARD OF MISSOURI BOTANICAL copyright reserved

GARDEN

Carcut planiflore var brevistyla.
C. brevistyla ABIn - Sched Le in Mich Feat flow Abyss. It, 79 Thei for seem to be well earth distinguished leg the short style, shick is equel tothe cray and skørter the the aprale, but there be'n no off god chamber fit the tyles varying i length itt othe for of C. plaifler I feel dele confide that that aught be until that Jolganghes Sperier. The original ones I find the lober of the corolla - fire expended in the top are the commet. Hes ben fin - Alaysin Schreepen II. 1486 on the Sinai, Botts, - Sonther Alexaniflens in Pensi. Ketsely sø Affghangt griffel 686. De ene The test Hooken fir a Thompson C. elegans Noë la 180 518. from the Zigni of Lan MISSOURI

BOTANICAL copyright reserved GARDEN



Casenta Curopaea Lui This will known and will chanden species extends from wintern lange throthy hotel Asia Cappany Lynn in the higher porter; of the country) to the Rimelaga Monatains. In southern IL is ful swed al llowing to the south of Hely Romandes seen speries for the right or Spin South of the Monney, or the north of Affrig - Al is fragin - Asii Minn, Carrapus, Versia, a & Shilet - the Mostly sude Montains grand Ance only I seems to have been the Method to their sports has give nice the good of the Method to Asquesto to regand to the presence or abrence of vales, but though I have exact a ranker of officiones, send to exhibit that organice. The my I have therefore day That it is always present Though in some sommelled their as to surrence defection of the southern their as to surrence of the strate so attack, itself its experience or build It is on the routing easily for I the dry corolle of all too much much much i pressing, it The speries C. Schkuhnaue Ofeif 1 the Synony falls to the ground? we few Halkand therefore the obronie only with its thich fleshy base a ette ned Colo, the their constle with obten Caria se the for their bified or trum Trales, the lange evan a loop participation of the stand of the convollant of the stand of the convollant top making difficulting to the stand of th MISSOURI

BOTANICAL copyright reserved cm GARDEN

If Syn: C. bruchystyla C. Koch L'ainaea XXII. 757

O. halophyta Fries nov. Mant. I. 8. 6 C. halophila Summ. Veg. Stand I. 191. 100 C. hyalina Boiss, in Sched. 13 C. Schkuhriann Opeif, bol. Jeit, Och 1845. for C. Segetum Asta - giorn bot ital A, 247 the form on asia 911 C. Miriae (Kindleg.) Dear. Blud. p. 27 Ja.C. Wilgaris Sens Syn I. 289. 1. C. Major Dach. Pin 219. De. Flor. Jr. III. DC produ. 1X, 452 8. C: monogyna Schmil. fl. Bohen et i herte 3 (tetrandra Moeach metter. 461 9 C. telvaspermen Jan - Hotel Job. C. tectocolora Mark. Lelia p. 215 18 Capillanis Edzeis, di Fransact XX 68 ej a more delig formett sa far form 14 ? Epitrifelyllen Donk Then Gerksty. 1844, 4804 4- C. Cpionistellin Buch, itw. 45 Cheropeen neforms Tries Lumm Veg. I. 191 16 Mely for portion (Moch Les 19. 1. 19 2. 6. fili formis of Law A. L. 507 Afres-chong Sillivis Cons MISSOURI BOTANICAL GARDER GEORGE ENGELMANN PAPERS

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

GARDEN

MISSOURI
BOTANICAL
GARDEN

Carlibus rapillacis, glomenlis bruter ovat. nate suffellis panolis pancis floris. floribus ante Sessil. bus plennyre 4-menis Calynis fore ad basis Strisi lobes ovato lamelle. acutis omminsonles, tuben correlae supentibes, laviai's ovato-lamedati, exectións. community

forme super capsula Vinflex is, taba fara long son lay. - Staminitus strye mallo brevionels, authoris paris sbrotunti, apindati, félaceto subveriontes, overie parle, reple medlo breven by, Le monte Jana, Kundstar, Kolschy R. aleja Kom. No 388. 6. sol nom. C. minor. n'herb. mulli, Kunstan I. Brant! Hell. Ken. A veg pearlianglant, sexully i textino I habit t. E. Comprer, Bot get dettet vilt stales even more slend. Distryed bythe Some of Caly x and Colle, Ste and Colon & Oplower I li log, seed lage - properto on the other hand the anti-lely a short styly
seent approach it to brevistyte, for said however it ei Aitraguelle lythe large flows with veg their lexture,


0 1 2 3 4 5 6 7 8 9 10

Copyright reserved

MISSOURI

BOTANICAL

GARDEN




Cascuta Versica De Risue - Herb Mus Par Conce filiforni, Storiter anti glowentif Zeni Ciber Courte s. orbiala seffelli, extendit about in. Calgini ompalette tobij ovater acuti: corollar tulem superati by, laviai's labole tile vix løngsviber exités, den peleli, ovatis absupte aven uti, dum papillozi, Han. mutto Greviontes. Igan. lannet files innunis fur volla acquait beg Styl bru. shalet over men men ples 207 MISSOURI

copyright reserved

BOTANICAL

GARDEN



Connett. Cpilinem Weihe, Archiv des Apoth. VIII. p. 54. (1824) Acichent. ir. E. 693. SC. Ond 1X. 452 C. densiflora, Søyer Willen Am. Soc. Linn. Paris I. p. 26. (1822)

C. densiflora, Søyer Willen Am. 201 (1826) C. major Riche 23, Col. only by rame, the descript is give IX p. 281 (1826). Epilinella as outoides, Pfeif, Boh. Jeil. och. 1845. Fask certist.


Des Moletas fludes sur les Censentes p. 64

elle Maure a 9 . 00 This well known and well chamber to species of the flaxfields of lumpe extends also into Asia to the Hol. bot. bol. Octorbay) Coppl (Brilit, Figari) & the Canaries (Hobb, Finley) and har ben volleil - the U. States (Manachusett, Oakes) similar lordities; on the banks of the Maine is half been seen on Wiria Faba (Roeden i He. Birkoff). - It is Indle der ornergin badia e it is pretele that Borkorphi Caggregate Hor intra I. p. 447 is their place " among flor of Bagel 2, while I tott tol. 9 - 2". - 2 have seen no indrien sperieur. The chamber solied on for a general separation for Carott are intable on are fred on wisher. The Colyx is Jeeply 5-lobe , beter les connète the i most alle peris bet mosto than i some amoin from 5 sepalles & The capsale is exactly constructed) as in the following species, the droup is complete, till a materialy it separates for the food of the affection edge still adhen, the app part of the representation The St, spind couries of 2 lawine with membrane but youth part are mailed into an off very this membrane but went to coges trave visione distint, as between the styles a triving portion of thee al make there up to paid downers of i some spens short, i other extress to the base of the caps les separates completely bearings an autor opening (a comment of a reports) of which the fighter consists) - which does not penetut it the rappele, The onifice beter the styles in Commonth, white The Dee w The interstyler opening the pertitute to the stiff of the stiff and by kill the stiffer of a send that the stiffer of almost the stiffer of the stiffer of almost the stiffer of Twice as leg as the etyle, (as that as that as base)

GARDEN

the Astigma the has the appearant of being clubskaper. Cercitis of the Cape of good Hope: while, the wey though not perfelled like all of theme; but month Cosily sessile, - i'l is probably a nature of Asia. If the struct of Me Sugar alpo Capalle politicals let ust It is with som besitating that Dear the species of the Read of the Come MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS The for I consider duplication dupostin Cathers MISSOURI BOTANICAL copyright reserved GARDEN

Consont, Kolschyane Briss Sing, or. I.7.29 Non C. Kolahyi Desmouling (= C. Exithyan van) Their species is debitegente by the large flows, with account bles of the corolla at with a lange at loose, regshagt Calex -It has been for by Kotchy in South Chrisis (N. 749). - A specie of collect by Sowiths i Armi the herowing the SI Peterts Agas such als be referre her, thought fly and & little Tomallen and the caly x mon



Cusoute plehella 1.1/1. floriber longier secontellati læxe fasori-leti, Celyvir carnosi labis ovatis acuti ptulo, corollae aquatio lænni rorolke telen sebæget by ovetri amling oulis, gralles papillosis, orenelatis, eretis 7. petalis, staminis brevinister, autheris oblogovorti frante subilete ægetibe, Igeni; augustis famme, ælgelle, fubritis in convitti, stiffiche its with overing out aprile subromin, agette, rapida brands. dem forme pan bæilen detir alle. met branch, in the Hook. Ken M' Alfghanston Griffith M. 688-691. on Alhaghi, Regamen, Letter? i Stein & flows equillely the plenty alex Bright red; inflowing losse, forflowed, flows all 1/3 his. Astragable for Mossichen by the much sunth flory with longe pushilly, the deeply In it rely, the stifffant ovate, not accumit lober of He molly the State Att Heib Host What Velens. I find two Theory one wolld ongull foller Huten Tota confirmate, thother toll. by. Sievers on the Alkai, which evilled belong here though they have nanver al men deeple doch Caley Cole, I satte mer avit, lober of It Corolla. Toles of bif. The form many be deliagules as vor affghang, the latter as var altaira, It is not impossible that MISSOURI

BOTANICAL

GARDEN

copyright reserved

cm

tt Catter maybe C. Asiatir of Vallar bat his own speril labeled thee it Hert. H.D. Veters. belons to C. planflore van Balayton MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS TO A DEPTH DESIGNATION OF THE SECOND STATES OF THE

copyright reserved

MISSOURI BOTANICAL GARDEN

Caro, pedsrellete Ledel, HAlfair. I. 293. Ic. C. 234 DC. Ord. 18. 453 excl. Syn. This is the smallest of these testing, flows sanly alin long frag stad the texterre; some small, knownte; skyler almost vonnet et beze separating tie the vije fruit fritte to mint opening, transcersely to the dissepied topped Capse de separation of the Committee in majoral of office of the separation of (720m) m/10m. I have example the offgen - Ime by Ledebour hundly a the Spens i his over Herbanian; At does not seent have be rolled by any subseget todavelle x H is som nenly alling to Combine the with both the speries of the scites with both the speries of the scites, and speries all styles. they it small flowing and the with opening of the Consule. MISSOURI

BOTANICAL copyright reserved GARDEN


Caesantes Stanberry Freder M. Aeg. p. 165, Chorisy Cour p. 175. +. 1. 1.2. Se Prod. 18. 453. (non Wight): Casset Arabira Desar, Et. 1.72 A ville make speries Vormon one i løgst, a lextulj uits Aubei. I har been collected - 13 yet by Bove 1954/ Auchen-Elog No 1418, de, Fischen Kneik, at strobble lefter ang om else by Lippi i Herb Vailland, ather it is mentions as Cascute sulphive olonis aggreptiona, flore niver et i Aubi Reippe solle the original sperior a scharge (N. 140) fe Lagain Thurs de distinguish to vaniti How it is possible to asonib to et option They are not see; nor it rapsale (barrele) exactly), as Desmonle Hetz The error however is easily around for as only the felly sign rapile will separa fritt bese with a very small opening mutite The phal for the Sinai Hongal (Anbria) i dityute by shorte perile Denn glomemler, conditi - sagittele auther, and Congre opper scales. He aggette Yvanity van aegyptière has long judvills satte storolon authers and maller, off bifið straler, which sometimes Reen to be crantity

0 1 2 3 4 5 6 7 8 9 10

Copyright reserved


GARDEN

MISSOURI
BOTANICAL
GARDEN



group fabrick while for it as all the otte allier by having a real bacrete capile which fells of out of the persolant Caley cutivity over by the closely eavedoping confla The Styles are subalete thech thish is usually week longer the A stigues.





C. capilal q. Mox6. C. rosea Jarquen out i Hert. M. 1880 ! This pretty spires form an exception for all the true Country by havi-the borrate Capsules, with fally off for the cally oping covered with the verience of the Corolla, leaving the. Jeen Stf relyx). Hei further distriguest by the sabelet. Myles with tops for a broad base, e. e. are tippe are stightedore only at the ope end j'/ for je Atte leaght. The Deeple Ind Colyx and could are veg minte cell, Almen Flats vy stort bearing an ovate or subglobore auther. Trale in the Com part of the tube of the corolle, not reach; to the mille, med on bifil dentale,

respect our with the Styles as la as one

much shorten the rapsole; rapsole rowned a

with the tale of the could, the laber of which on comment of log begge intre How our 1/2 le 2006. ilong slight venuerose. kudrå grad ela len - Crotalaris junea (Rox6 / For Artemsia of Himalaya Glegend N/1550, - On Mynn, Hardeya 7-10,000 feet by, & Hough _ in the Kens On small herbes in Kunning 12.000 fut ligh Shackney & westerbottom, M3. - 14. Men Oding MISSOURI

BOTANICAL copyright reserved GARDEN

GARDEN

cm